

Monoclonal Antibodies in Nephrotic Syndrome

Gian Marco Ghiggeri

Andrea Angeletti

Istituto Giannina Gaslini, Genova



Monoclonal Antibodies (-MAB)

Mono



**SINGLE
Or
ONE**

- clonal




**IDENTICAL/SAME
GROUP OF CELLS
DERIVED FROM SINGLE
PARENT CELLS**

Antibody


**ANTIBODY PRODUCED BY
ONE
TYPE OF IMMUNE CELLS**

- ANTIBODY BINDS TO SINGLE SPECIFIC ANTIGENS (TARGET)
- HIGHLY SPECIFIC
- UNIFORM
- LARGE QUANTITIES



(Chinese hamster ovary)

(murine myeloma)




Alta

IMMUNOGENICITÀ

Bassa

**Primary
NS**

**Secondary
NS**



Nephrotic Syndrome: Definition



Steroid Sensitive
80%

Steroid resistant
20%

↓

Steroid Dependent
50%

↙ ↘

Multidrug Dependent
(MMF/Tac+Ste)

Multidrug Resistant

↓

**Dialysis/
Transplantation**
50%

Historical use of Anti-CD20

- Introduced in the late '90s to treat NH lymphoma
- Used to treat autoimmune disease with predominant humoral component / pathogenetic auto-ab
- *Found to reduce proteinuria in children with nephrotic syndrome secondary to PTLD (Nozu, Pediatric Nephrol 2005; Pescovitz, NEJM 2006)*

Historical use of Anti-CD20

Fig. 1 Renal biopsy 2 months after transplantation (a, b) and 3 years after transplantation (c). **a** PAS staining shows minor glomerular abnormalities. **b** Electron microscopy indicates 60% foot process fusion. **c** No foot process fusion is detectable





Fig. 2 Clinical course in terms of peripheral B cell counts, serum IgG levels, and ratio of urinary protein to urinary creatinine (U-P/U-Cr). **a** Peripheral B cells disappeared immediately after the rituximab treatment, while serum IgG levels decreased gradually. **b** Urinary protein decreased immediately after the rituximab administration



Nephrotic Syndrome: Most Relevant RCTs for RITUXIMAB

Table 1

Major published clinical studies testing biologics in pediatric MCD and FSGS

Reference	Disease	N	F/U (mo) ^a	Study design	Dose	CR (%)	Comments
Rituximab Ravani <i>et al.</i> [40]	SDNS	30	22 (1–60)	RCT	375 mg/m ² (1–4 doses) vs. no therapy	N/A	Add-on RTX did not significantly increase the rate of remission after steroid tapering
Iijima <i>et al.</i> [32]	FRNS/SDNS	48	12	RCT	375 mg/m ² (4 doses) vs. placebo	81	Mean daily steroid dose after randomization was significantly lower in the RTX group than in the placebo group
Ruggenenti <i>et al.</i> [39]	SDNS	30	12	Off-on trial	375 mg/m ² (1–2 doses)	100	Patients were in remission at treatment start. During the 24 mo after RTX administration, patients experienced fewer recurrences than in the 24 mo prior
Ravani <i>et al.</i> [42]	multidrug dependent NS	46	6	Off-on trial	375 mg/m ² (1–5 doses)	48	At 18 mo, 20% of children were still in prednisone and CNI-free remission
Sinha <i>et al.</i> (2015)	multidrug dependent NS	19 5	12	Prospective cohort	375 mg/m ² (2–4 doses)	38	Remission rate was 57.6% in steroid dependent, 33% in CNI-dependent and 12% in steroid + CNI-resistant patients at 12 months
Sellier-Lecler <i>et al.</i> [33]	SDNS	30	38 (25–51)	Prospective cohort	375 mg/m ² (1–4 doses)	63	19 patients underwent remission without steroids. 11 patients relapsed, ten of them received RTX re-treatment
Fujinaga <i>et al.</i> [31]	SDNS	10	17 (13–21)	Prospective cohort	375 mg/m ² (1 dose)	N/A	All patients had MCD
Kamei <i>et al.</i> [37]	SDNS	12	12	Prospective cohort	375 mg/m ² (1 dose)	25	After steroid withdrawal, 9 patients relapsed, but 3 remained in remission without steroids for >1 year

Nephrotic Syndrome: Most Relevant RCTs for RITUXIMAB

TRY RESCUE 2

Ghiggeri, cJASN 2011,
Ghiggeri, JASN 2015

RCRNS

Iijima, Lancet 2014

RITURNS

Basu, JAMAP 2018


OFA 2

Ghiggeri, JASN 2021

RTX 4

Ghiggeri, end of recruitment

Long-Term Efficacy and Safety of Repeated Rituximab to Maintain Remission in Idiopathic Childhood Nephrotic Syndrome: An International Study



Children receiving repeated courses of rituximab for FRSDNS experience an improving clinical response. Side effects appear acceptable


Number at risk

Course 1	49	14	1	0	0	0	0	0
Course 2	49	18	6	1	0	0	0	0
Course 3	49	24	9	4	0	0	0	0
Course 4	49	40	14	6	4	3	1	0
	0	12	24	36	48	60	72	84

Time from RTX (months)

(Chinese hamster ovary)

(murine myeloma)



Alta

IMMUNOGENICITÀ

Bassa

RTX vs. OFA

Human or Chimeric Monoclonal Anti-CD20 Antibodies for Children with Nephrotic Syndrome: A Superiority Randomized Controlled Trial


JASN[®]
JOURNAL OF THE AMERICAN SOCIETY OF NEPHROLOGY

METHODS

Open-label, two-parallel-arm, randomized controlled trial



140 patients (2-24yrs) with Steroid Dependent NS




Endpoints

Relapse at 12 and 24 months

OUTCOME

Relapse

	OFA	RTX
12 months	37/70	36/70
24 months	53/70	46/70




Safety

No relevant AEs

Conclusion

Ofatumumab was not superior to rituximab in maintaining remission in children and young adults with steroid- and calcineurin inhibitor-dependent nephrotic syndrome.


doi: 10.1681/ASN.2021040561



Subject at Risk

Months	0	6	12	18	24
OFA	70	57	33	20	17
RTX	70	52	34	29	24


≤9 years



Subjects at Risk

Months	0	6	12	18	24
OFA _{≤9 yrs}	32	25	12	8	5
RTX _{≤9 yrs}	31	25	14	13	11

>9 years







Subjects at Risk

Months	0	6	12	18	24
OFA _{>9 yrs.}	38	32	21	12	12
RTX _{>9 yrs}	39	27	20	17	13

Safety


- Used in one million of patients with hematologic malignancies (first line / maintenance)
- First **infusion adverse reactions** (bronchospasm, cough, chills, rash, fever, headache); mild / absent thereafter
- Systematic review in RCTs of RA treated with biologic therapy (N = 29,423): **no increased risk of malignancies** (Lopez-Olivo JAMA 2012)
- 23 cases of **PML** in > 500,000 patients; had either B-cell cancer (20) or lupus (3); all taking chemotherapy - **no cases in NS**

Safety




—●— Ofatumumab —▲— Rituximab

Safety



Circulating Anti-Rituximab Antibodies Do Not Affect Response to Rituximab in Steroid-Dependent Nephrotic Syndrome




Subjects at Risk

Months	0	6	12	18	24
anti-RTXpos	14	11	10	9	8
anti-RTX neg	40	36	30	27	22

Key points

- RTX is useful in the treatment of some forms of INS in children (combined steroid/CNI dependent forms)
- RTX high superior to PDN on pure dependent forms
- OFatumumab is equal to Rituximab
- RTX ineffective in multidrug-resistant INS

Nephrotic Syndrome: Definition




Nephrotic Syndrome: Definition

Steroid Sensitive
80%



Steroid Dependent
50%



STEROID/MULTIDRUG RESISTANT NS

Reference	Disease	N	Study	Dose	CR (%)
Rituximab					
Bagga A, <i>et al.</i> , 2007 (20)	SRNS	5	Case series	375mg/m ² (4 doses)	60
Gulati A, <i>et al.</i> , 2010 (21)	SRNS	33	Multicentric Cohort Study	375mg/m ² (4 doses)	27
Magnasco A, <i>et al.</i> , 2012 (22)	SRNS	31	RCT	375mg/m ² (2 doses) vs 375mg/m ² + Cyc	30
Ofatumumab					
Basu B, <i>et al.</i> , 2014 (49)	multidrug resistant NS	5	Case series	300mg/m ² followed by 5 weekly infusions (2g/m ²)	80
Bonanni A, (50) <i>et al.</i> , 2015	multidrug resistant NS	4	Case series	375-700 mg/m ² (1 dose)	25
Wang CS, <i>et al.</i> , 2017 (52)	SRNS	4	Case series	300mg/m ² followed by 5 weekly infusions (2g/m ²)	75
Ravani P, <i>et al.</i> , 2020 (53)	multidrug resistant NS	7	RCT	1,500mg/m ² (1 dose)	0

STEROID/MULTIDRUG RESISTANT NS

- **Obinutuzumab** (lesson from Lupus nephritis)
- **Anti-CD38 Abs +Rituximab** (trial in development)
- Finerenone
- SGLT2i

A Study to Evaluate the Efficacy and Safety of Obinutuzumab Versus MMF in Participants With Childhood Onset Idiopathic Nephrotic Syndrome (INShore)

A Phase 3 study

Study Type :	Interventional (Clinical Trial)
Estimated Enrollment :	80 participants
Allocation:	Randomized
Intervention Model:	Parallel Assignment
Masking:	None (Open Label)
Primary Purpose:	Treatment
Actual Study Start Date :	March 29, 2023
Estimated Primary Completion Date :	December 1, 2025
Estimated Study Completion Date :	August 15, 2026

1) Title

Efficacy and tolerance of the association between chimeric monoclonal anti-CD20 (Rituximab Biosimilar) and monoclonal anti-CD38 (Daratumumab) antibodies in the treatment of childhood multidrug dependent and resistant and in post transplant recurrence of nephrotic syndrome.

A Phase 2 study (the DUAL1)

2) Trial Registration

NCT05704400

Eudract: 2022-001769-11

ENROLLING PHASE

BIOMARKERS-PATHOGENESIS


- CD20
- Memory/IgM CD20 (in progress)
- Antibodies (in developement)
- Serum/Urinary proteins (in progress)

MOMA* Project

* Multi Omics


Molecular Analysis

Nephrotic Syndrome




PLASMA

Name	Total
Lipidi_C18_Pos_plasma_D_014	970
Lipidi_C18_Pos_plasma_D_015	971
Lipidi_C18_Pos_plasma_D_021	968
Lipidi_C18_Pos_plasma_IPA_017	965
Lipidi_C18_Pos_plasma_IPA_020	960
Lipidi_C18_Pos_plasma_IPA_024	963
Lipidi_C18_Pos_plasma_MANN_016	959
Lipidi_C18_Pos_plasma_MANN_019	960
Lipidi_C18_Pos_plasma_MANN_022	959






URINE

Name	Total
Lipidi_C18_Pos_urine_D_044	248
Lipidi_C18_Pos_urine_D_051	248
Lipidi_C18_Pos_urine_D_053	248
Lipidi_C18_Pos_urine_IPA_045	241
Lipidi_C18_Pos_urine_IPA_048	243
Lipidi_C18_Pos_urine_IPA_050	243
Lipidi_C18_Pos_urine_MANN_...	245
Lipidi_C18_Pos_urine_MANN_...	245
Lipidi_C18_Pos_urine_MANN_...	246




Weighted Gene Co-Expression Network Analysis (WGCNA)

(Langfelder P and Horvath S, BMC Bioinformatics, 2008)



Machine learning and supervised cluster analysis



unpublished data



Lab Crew

Francesca Lugani

Maurizio Bruschi

Gianluca Caridi

Giovanni Candiano

Xhuliana Kasana

Sonia Spinelli

Andrea Garbarino



Dr. Enrico Verrina

andreaangeletti@gaslini.org

gmarcoghiggeri@gaslini.org

+39.3284140532





In memory of Prof R Gusmano

Thanks to Dr G Ghiggeri